

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 2007		2. REPORT TYPE		3. DATES COVERED 00-00-2007 to 00-00-2007	
4. TITLE AND SUBTITLE Polar Clouds from Space Shuttle Exhaust				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory, 4555 Overlook Avenue SW, Washington, DC, 20375				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 2	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Polar Clouds from Space Shuttle Exhaust

M.H. Stevens and C.R. Englert
Space Science Division

Introduction: The Earth's mesosphere lies above the stratosphere in the region between 50 and 90 km altitude, far above where airplanes or balloons can fly. In the unique polar summer mesosphere, the temperature plummets below 150 K (-190°F), making this region the coldest on Earth, so inaccessible that it is sometimes like studying the atmosphere of another planet. In this extremely rarefied and dry place, water ice particles are found in narrow layers near 82 km altitude called polar mesospheric clouds (PMCs). PMCs are not known in the published record until 1885.

The specific processes leading to the formation of PMCs are disputed. However, some evidence indicates that they became brighter and more frequent in the late 20th century, leading some scientists to argue that they are indicators of global climate change. One hypothesis reasons that increasing amounts of methane (CH_4) emitted at the Earth's surface by industrial and agricultural processes increase the humidity of the upper atmosphere as the methane is broken down by ultraviolet sunlight to form water vapor.

Recently, NRL's Space Science Division scientists have complicated this hypothesis by identifying a new source for PMCs, challenging long-held beliefs about the meteorology of the upper atmosphere. Using satellite observations from NRL's Middle Atmosphere High Resolution Spectrograph Investigation (MAHRSI), Stevens et al.¹ showed that the exhaust plume from the space shuttle can be transported all the way from the east coast of the United States to the Arctic summer mesosphere to form PMCs. Additional data from other experiments now reveal that this phenomenon has occurred over both poles. Here we present the initial observations leading to the discovery, and discuss its scientific impact.

The Discovery: MAHRSI was an experiment designed to measure hydroxyl (OH) in the Earth's mesosphere on a satellite deployed and retrieved by the crew of the space shuttle. It flew on two one-week missions in November 1994 (STS-66) and August 1997 (STS-85). MAHRSI observed near-ultraviolet light ($\sim 309\text{ nm}$) from the atmosphere at high spectral resolution (0.02 nm), which allowed for the discrimination of OH solar fluorescence from the bright Rayleigh scattered background. PMCs were never considered as a science objective of MAHRSI, but were unexpectedly detected while analyzing data at the Kennedy Space Center during the late summer mission in 1997.²

OH is created when water vapor is destroyed by solar ultraviolet radiation and it is therefore sometimes used as a proxy for water vapor. Water vapor is the primary effluent of the shuttle's main engines and is copiously injected in a relatively narrow layer between 105 and 115 km altitude off the east coast of the United States as the shuttle accelerates to orbit.


As shown in Figs. 3(a) and 3(b), one to two days after the launch of STS-85, unexpectedly bright OH intensities were observed by MAHRSI in the upper mesosphere at Arctic latitudes (circled in red). Figure 3(c) shows that about a week after launch, PMCs were also observed over much of northern North America by MAHRSI, as indicated by the red crosses. Using observations from a German companion experiment called Cryogenic Infrared Spectrometers and Telescopes for the Atmosphere (CRISTA), Stevens et al.¹ found that the water content of these clouds ($\sim 100\text{ t}$) was consistent with the amount available in the shuttle plume ($\sim 300\text{ t}$). This was the first evidence that a launch vehicle plume injected in the subtropics could create Arctic PMCs. To our knowledge, the global scale transport of this exhaust plume had never been considered before.

The Scientific Impact: We have since assembled more data from a variety of ground-based and satellite experiments. For example, we found that another shuttle exhaust plume (STS-107) created polar clouds, but this time they were in the Antarctic summer, contributing 10-20% to the total ice content of PMCs observed that season.³ Proposed trends in cloud brightness since 1980 are not larger than this. This calls into question the interpretation of late 20th-century PMC trends solely in the context of global climate change.

The Upper Atmospheric Physics Branch at NRL's Space Science Division is poised to build on the success of MAHRSI with a new instrument called the Spatial Heterodyne IMager for MESospheric Radicals (SHIMMER). This instrument uses innovative technology called spatial heterodyne spectroscopy that allows for higher spectral resolution and throughput while minimizing mass and volume. SHIMMER is scheduled to launch in 2007 on a much longer, one-year mission to measure OH in the middle atmosphere.

Acknowledgments: The authors are grateful to Jörg Gumbel, Klaus Grossmann, Paul Hartogh, Markus Rapp, Robert Meier, Xinzhaoh Chu, Matthew DeLand, and John Plane for providing data and model results in valuable support of this work. We are also grateful to David E. Siskind for many productive discussions and suggestions.

[Sponsored by ONR and NASA]

**FIGURE 3**

STS-85 carried the Middle Atmosphere High Resolution Spectrograph Investigation (MAHRSI) into orbit in August 1997. MAHRSI observed the shuttle's exhaust plume in the Arctic (circled in red) by measuring solar resonance fluorescence of OH, where 80–85 km radiances are referenced to the color bar. MAHRSI scans the limb of the Earth, so that OH observed viewing toward 80–85 km can originate at altitudes higher than this along the line of sight. A week after launch, MAHRSI observed polar mesospheric clouds over northern North America (red crosses, Fig. 3(c)). The amount of water in the observed PMCs is consistent with the amount injected by the shuttle.

References

- ¹ M.H. Stevens, J. Gumbel, C.R. Englert, K.U. Grossmann, M. Rapp, and P. Hartogh, "Polar Mesospheric Clouds Formed from Space Shuttle Exhaust," *Geophys. Res. Lett.* **30**(10), 1546 (2003). doi: 10.1029/2003GL017249
- ² M.H. Stevens, "Heavenly Harbingers," *Smithsonian*, November, 2001, 20-23.
- ³ M.H. Stevens, R.R. Meier, X. Chu, M.T. DeLand, and J.M.C. Plane, "Antarctic Mesospheric Clouds Formed from Space Shuttle Exhaust," *Geophys. Res. Lett.* **32**, L13810 (2005). doi: 10.1029/2005GL023054